arXiv:0803.3340v2 [math.OA] 25 Mar 2008

Type III_1 factors generated by regular representations of infinite dimensional nilpotent group $B_0^{\mathbb{N}}$

Alexandre Kosyak

Max-Planck-Institut für Mathematik, Vivatsgasse 7, D-53111 Bonn, Germany

Institute of Mathematics, Ukrainian National Academy of Sciences, 3
Tereshchenkivs'ka, Kyiv, 01601, Ukraine,
E-mail: kosyak01@yahoo.com, kosyak@imath.kiev.ua
tel.: 38044 2346153 (office), 38044 5656758 (home), fax: 38044 2352010

Abstract

We study the von Neumann algebra, generated by the unitary representations of infinite-dimensional groups nilpotent group $B_0^{\mathbb{N}}$. The conditions of the irreducibility of the regular and quasiregular representations of infinite-dimensional groups (associated with some quasi-invariant measures) are given by the so-called Ismagilov conjecture (see [1,2,9,10,11]). In this case the corresponding von Neumann algebra is type I_∞ factor. When the regular representation is reducible we find the sufficient conditions on the measure for the von Neumann algebra to be factor (see [13,14]). In the present article we determine the type of corresponding factors. Namely we prove that the von Neumann algebra generated by the regular representations of infinite-dimensional nilpotent group $B_0^{\mathbb{N}}$ is type III_1 hyperfinite factor. The case of the nilpotent group $B_0^{\mathbb{Z}}$ of infinite in both directions matrices will be studied in [6].

Key words: von Neumann algebra, type III_1 factor, unitary representation, infinite-dimensional groups, nilpotent groups, regular representations, irreducibility, infinite tensor products, Gaussian measures, Ismagilov conjecture
1991 MSC: 22E65, (28D25, 17B65, 28C20)

⋆ The author would like to thank the Max-Planck-Institute of Mathematics, Bonn for the hospitality. The partial financial support by the DFG project 436 UKR 113/87 is gratefully acknowledged.

* Corresponding author
Email address: `kosyak01@yahoo.com, kosyak@imath.kiev.ua` (Alexandre Kosyak).

Contents

1 Regular representations 3

2 Von Neuman algebras generated by the regular representations 5

3 Type III_1 factor 6

4 The von Neumann algebra M_ϕ is trivial 13

5 Example of the measure 21

6 Modular operator 22

7 The uniqueness of the constructed factor 24

References 25

1 Regular representations

Let us consider the group $\tilde{G} = B^{\mathbb{N}}$ of all upper-triangular real matrices of infinite order with unities on the diagonal

$$\tilde{G} = B^{\mathbb{N}} = \{I + x \mid x = \sum_{1 \leq k < n} x_{kn} E_{kn}\},$$

and its subgroup

$$G = B_0^{\mathbb{N}} = \{I + x \in B^{\mathbb{N}} \mid \ x \text{ is finite}\},$$

where E_{kn} is an infinite-dimensional matrix with 1 at the place $k, n \in \mathbb{N}$ and zeros elsewhere, $x = (x_{kn})_{k<n}$ is *finite* means that $x_{kn} = 0$ for all (k, n) except for a finite number of indices $k, n \in \mathbb{N}$.

Obviously, $B_0^{\mathbb{N}} = \varinjlim_n B(n, \mathbb{R})$ is the inductive limit of the group $B(n, \mathbb{R})$ of real upper-triangular matrices with units on the principal diagonal

$$B(n, \mathbb{R}) = \{I + \sum_{1 \leq k < r \leq n} x_{kr} E_{kr} \mid x_{kr} \in \mathbb{R}\}$$

with respect to the imbedding $B(n, \mathbb{R}) \ni x \mapsto x + E_{n+1n+1} \in B(n + 1, \mathbb{R})$.

We define the Gaussian measure μ_b on the group $B^{\mathbb{N}}$ in the following way

$$d\mu_b(x) = \otimes_{1 \leq k < n} (b_{kn}/\pi)^{1/2} \exp(-b_{kn} x_{kn}^2) dx_{kn} = \otimes_{k<n} d\mu_{b_{kn}}(x_{kn}), \qquad (1)$$

where $b = (b_{kn})_{k<n}$ is some set of positive numbers.

Let us denote by R and L the right and the left action of the group $B^{\mathbb{N}}$ on itself: $R_s(t) = ts^{-1},\ \ L_s(t) = st,\ \ s, t \in B^{\mathbb{N}}$ and by $\Phi : B^{\mathbb{N}} \mapsto B^{\mathbb{N}},\ \Phi(I + x) := (I + x)^{-1}$ the inverse mapping. It is known [9,10] that

Lemma 1 *$\mu_b^{R_t} \sim \mu_b\ \forall t \in B_0^{\mathbb{N}}$ for any set $b = (b_{kn})_{k<n}$.*

Lemma 2 *$\mu_b^{L_t} \sim \mu_b\ \forall t \in B_0^{\mathbb{N}}$ if and only if $S_{kn}^L(b) < \infty,\ \forall k < n$, where*

$$S_{kn}^L(b) = \sum_{m=n+1}^{\infty} \frac{b_{km}}{b_{nm}}.$$

Lemma 3 *$\mu_b^{L_t} \perp \mu_b\ \forall t \in B_0^{\mathbb{N}} \backslash \{e\} \Leftrightarrow S_{kn}^L(b) = \infty\ \ \forall k < n.$*

Lemma 4 [12] *If $E(b) = \sum_{k<n} S_{kn}^L(b)(b_{kn})^{-1} < \infty$, then $\mu_b^{\Phi} \sim \mu_b$.*

Lemma 5 [12] *The measure μ_b on $B^{\mathbb{N}}$ is $B_0^{\mathbb{N}}$ ergodic with respect to the right action.*

Let $\alpha : G \to \mathrm{Aut}(X)$ be a measurable action of a group G on the measurable space X. We recall that a measure μ on the space X is G-ergodic if $f(\alpha_t(x)) = f(x)\ \forall t \in G$ implies $f(x) = const\ \mu$ a.e. for all functions $f \in L^1(X, \mu)$.

Remark 6 [13] *If $\mu_b^{\Phi} \sim \mu_b$ then $\mu_b^{L_t} \sim \mu_b\ \forall t \in B_0^{\mathbb{N}}$.*

PROOF. This follows from the fact that the inversion Φ replace the right and the left action: $R_t \circ \Phi = \Phi \circ L_t\ \forall t \in B^{\mathbb{N}}$. Indeed, if we denote $\mu^f(\cdot) = \mu(f^{-1}(\cdot))$ we have $(\mu^f)^g = \mu^{f \circ g}$. Hence

$$\mu_b \sim \mu_b^{R_t} \sim (\mu_b^{R_t})^{\Phi} = \mu_b^{R_t \circ \Phi} = \mu_b^{\Phi \circ L_t} = (\mu_b^{\Phi})^{L_t} \sim \mu_b^{L_t}.$$

□

If $\mu_b^{R_t} \sim \mu_b$ and $\mu_b^{L_t} \sim \mu_b\ \forall t \in B_0^{\mathbb{N}}$, one can define in a natural way (see [9,10]), an analogue of the right $T^{R,b}$ and left $T^{L,b}$ representation of the group $B_0^{\mathbb{N}}$ in Hilbert space $H_b = L_2(B^{\mathbb{N}}, d\mu_b)$

$$T^{R,b},\ T^{L,b} : B_0^{\mathbb{N}} \to U(H_b = L_2(B^{\mathbb{N}}, d\mu_b)),$$

$$(T_t^{R,b} f)(x) = (d\mu_b(xt)/d\mu_b(x))^{1/2} f(xt),$$

$$(T_s^{L,b} f)(x) = (d\mu_b(s^{-1}x)/d\mu_b(x))^{1/2} f(s^{-1}x).$$

2 Von Neuman algebras generated by the regular representations

Let $\mathfrak{A}^{R,b} = (T_t^{R,b} \mid t \in B_0^{\mathbb{N}})''$ (resp. $\mathfrak{A}^{L,b} = (T_s^{L,b} \mid s \in B_0^{\mathbb{N}})''$) be the von Neumann algebras generated by the right $T^{R,b}$ (resp. the left $T^{L,b}$) regular representation of the group $B_0^{\mathbb{N}}$.

Theorem 7 [12] *If $E(b) < \infty$ then $\mu_b^{\Phi} \sim \mu_b$. In this case the left regular representation is well defined and the commutation theorem holds:*

$$(\mathfrak{A}^{R,b})' = \mathfrak{A}^{L,b}. \tag{2}$$

Moreover, the operator J_{μ_b} given by

$$(J_{\mu_b} f)(x) = (d\mu_b(x^{-1})/d\mu_b(x))^{1/2}\overline{f(x^{-1})} \tag{3}$$

is an intertwining operator:

$$T_t^{L,b} = J_{\mu_b} T_t^{R,b} J_{\mu_b},\ t \in B_0^{\mathbb{N}} \quad \text{and } J_{\mu_b} \mathfrak{A}^{R,b} J_{\mu_b} = \mathfrak{A}^{L,b}.$$

If $\mu_b^{L_t} \perp \mu_b \ \forall t \in B_0^{\mathbb{N}} \backslash \{e\}$ one can't define the left regular representation of the group $B_0^{\mathbb{N}}$. Moreover the following theorem holds

Theorem 8 *The right regular representation $T^{R,b} : B_0^{\mathbb{N}} \mapsto U(H_b)$ is irreducible if and only if $\mu_b^{L_s} \perp \mu_b \ \forall s \in B_0^{\mathbb{N}} \backslash \{0\}$.*

Corollary 9 *The von Neumann algebra $\mathfrak{A}^{R,b}$ is a type I_∞ factor if*

$\mu_b^{L_s} \perp \mu_b \ \forall s \in B_0^{\mathbb{N}} \backslash \{0\}$.

Let us assume now that $\mu_b^{L_t} \sim \mu_b \ \forall t \in B_0^{\mathbb{N}} \backslash \{e\}$. In this case the right regular representation and the left regular representation of the group $B_0^{\mathbb{N}}$ are well defined.

In [13] the condition were studied *when the von Neumann algebra $\mathfrak{A}^{R,b}$ is factor*, i.e.

$$\mathfrak{A}^{R,b} \cap (\mathfrak{A}^{R,b})' = \{\lambda \mathbf{I} | \lambda \in \mathbb{C}^1\}.$$

Since $T_t^{L,b} \in (\mathfrak{A}^{R,b})' \ \forall t \in B_0^{\mathbb{N}}$, we have $\mathfrak{A}^{L,b} \subset (\mathfrak{A}^{R,b})'$, hence

$$\mathfrak{A}^{R,b} \cap (\mathfrak{A}^{R,b})' \subset (\mathfrak{A}^{L,b})' \cap (\mathfrak{A}^{R,b})' = (\mathfrak{A}^{R,b} \cup \mathfrak{A}^{L,b})'. \tag{4}$$

The last relation shows that $\mathfrak{A}^{R,b}$ is factor if the representation

$$B_0^{\mathbb{N}} \times B_0^{\mathbb{N}} \ni (t,\, s) \to T_t^{R,b} T_s^{L,b} \in U(H_b)$$

is irreducible.

Let us denote by $\mathfrak{A}^{R,L,b}$ the the von Neumann algebras generated by the right $T^{R,b}$ and the left $T^{L,b}$ regular representations of the group $B_0^{\mathbb{N}}$:

$$\mathfrak{A}^{R,L,b} = (T_t^{R,b}, T_s^{L,b} \mid t, s \in B_0^{\mathbb{N}})'' = (\mathfrak{A}^{R,b} \cup \mathfrak{A}^{L,b})''.$$

Let us denote

$$S_{kn}^{R,L}(b) = \sum_{m=n+1}^{\infty} \frac{b_{km}}{S_{nm}^{L}(b)}, \ \ k < n.$$

Theorem 10 [13] *The representation*

$$B_0^{\mathbb{N}} \times B_0^{\mathbb{N}} \ni (t,\ s) \to T_t^{R,b} T_s^{L,b} \in U(H_b)$$

is irreducible if $S_{kn}^{R,L}(b) = \infty, \ \forall k < n.$

Corollary 11 *The von Neumann algebra* $\mathfrak{A}^{R,b}$ *is factor if* $S_{kn}^{R,L}(b) = \infty \, \forall k < n.$

3 Type III_1 factor

Let us denote as before $M = \mathfrak{A}^{L,b} = (T_s^{L,b} \mid s \in B_0^{\mathbb{N}})''$, $\mathfrak{A}^{R,b} = (T_t^{R,b} \mid t \in B_0^{\mathbb{N}})''$.

Theorem 12 *If* $S_{kn}^{R,L}(b) = \infty, \ \forall k < n$ *then the von Neumann algebra* $\mathfrak{A}^{L,b}$ *(and hence* $\mathfrak{A}^{R,b}$*) is* III_1 *factor.*

PROOF. The proof is based on Lemma 13 and 14, we shall prove them later.

Using (3) we conclude that the modular operator Δ is defined as follows

$$(\Delta f)(x) = (d\mu_b(x)/d\mu_b(x^{-1}))f(x). \tag{5}$$

Lemma 13 *We have*

$$Sp\Delta = [0, \infty).$$

We have $Sp\Delta\phi = Sp\Delta = [0, \infty)$, where $\phi(a) = (a\mathbf{1}, \mathbf{1})_{H_b}, \ a \in M = \mathfrak{A}^{L,b}$. The centralizer M_ϕ of ϕ is defined by the equality

$$M_\phi = \{a \in M \mid \sigma_t^\phi(a) \ \forall t \in \mathbb{R}\}$$

where $\sigma_t^\phi(a) = \Delta^{it} a \Delta^{-it}$. For every projection $e \neq 0, e \in M_\phi$, a faithful semifinite normal weight ϕ_e on the reduced von Neumann algebra $eMe =$

$\{a \in M;\ ea = ae = a\}$ is defined by the equality

$$\phi_e(a) = \phi(a)\ \ \forall a \in eMe, a \geq 0.$$

One has the formula

$$S(M) = \bigcap_{e \neq 0} Sp\Delta_{\phi_e}, \tag{6}$$

where e varies over the nonzero projection of M_ϕ (see[4] p.472).

Lemma 14 *The von Neumann algebra M_ϕ is trivial.*

In this case

$$S(M) = Sp\Delta = [0, \infty),$$

so the von Neumann algebra $\mathfrak{A}^{L,b}$ (and hence algebra $\mathfrak{A}^{R,b}$) is type III$_1$ factor. $\square$

Proof of Lemma 14. We show that

$$M_\phi = (\Delta^{it},\, T_s^{R,b} \mid t \in \mathbb{R},\, s \in B_0^{\mathbb{N}})'. \tag{7}$$

So M_ϕ is trivial means that the set of operators

$$(\Delta^{it},\, T_s^{R,b} \mid t \in \mathbb{R},\, s \in B_0^{\mathbb{N}}) \tag{8}$$

is **irreducible**. To prove (7) we get

$$M_\phi = (a \in \mathfrak{A}^{L,b} \mid \Delta^{it} a = a\Delta^{it},\ \forall t \in \mathbb{R}) = (\Delta^{it} \mid t \in \mathbb{R})' \cap \mathfrak{A}^{L,b}$$

$$= (\Delta^{it} \mid t \in \mathbb{R})' \cap (\mathfrak{A}^{R,b})' = (\Delta^{it} \mid t \in \mathbb{R})' \cap (T_s^{R,b} \mid s \in B_0^{\mathbb{N}})' =$$
$$(\Delta^{it},\, T_s^{R,b} \mid t \in \mathbb{R},\, s \in B_0^{\mathbb{N}})'$$

Definition. Recall (c.f. e.g. [5]) that a non necessarily bounded self-adjoint operator A in a Hilbert space H is said to be *affiliated* with a von Neumann algebra M of operators in this Hilbert space H, if $\exp(itA) \in M$ for all $t \in \mathbb{R}$. One then writes $A\ \eta\ M$.

To prove the irreducibility of $(\Delta^{it},\, T_s^{R,b} \mid t \in \mathbb{R},\, s \in B_0^{\mathbb{N}})$ it is sufficient to prove (see [10] p.258) that operators $f(x) \mapsto x_{kn} f(x)$ of multiplication in the space H_b by the independent variables x_{kn} are affiliated to the von Neumann algebra

$$(M_\phi)' = (\Delta^{it},\, T_s^{R,b} \mid t \in \mathbb{R},\, s \in B_0^{\mathbb{N}})''.$$

In this case the operator A commuting with Δ^{it} and $T_s^{R,b}$ is operator of multiplication by some function $a(x)$. If we use commutation relation $[A, T_s^{R,b}] = 0,\ s \in B_0^{\mathbb{N}}$ we obtain $a(x) = a(xs) \bmod \mu$. Using the ergodocity of the measure μ_b with respect of the right action of the group $B_0^{\mathbb{N}}$ we conclude that $a(x) = \text{const} \bmod \mu$ i.e. A is scalar operator.

If we denote

$$A_{kn}^R=(d/dt)T_{I+tE_{kn}}^{R,b}\mid_{t=o}$$

we have (see for example [9,10,11])

$$A_{kn}^R=\sum_{r=1}^{k-1}x_{kr}D_{rn}+D_{kn},\quad 1\leq k<n. \tag{9}$$

The direct calculation shows that

$$[A_{13}^R,[A_{23}^R,\,\ln\Delta]]=2b_{13}x_{12}, \tag{10}$$

$$[A_{12}^R,[A_{23}^R,\,\ln\Delta]]=2b_{13}x_{13}. \tag{11}$$

Idea: to obtain in a similar way all variables x_{kn}.

Let us denote by X^{-1} the inverse matrix to the upper triangular matrix $X=I+x=I+\sum_{k<n}x_{kn}E_{kn}\in B^{\mathbb{N}}$

$$X^{-1}=(I+x)^{-1}=I+\sum_{k<n}x_{kn}^{-1}E_{kn}\in B^{\mathbb{N}}.$$

We have by definition $X^{-1}X=XX^{-1}=I$ hence

$$\Big(XX^{-1}\Big)_{kn}=\sum_{r=k}^{n}x_{kr}x_{rn}^{-1}=\delta_{kn}=\sum_{r=k}^{n}x_{kr}^{-1}x_{rn}=\Big(X^{-1}X\Big)_{kn},\quad k\leq n, \tag{12}$$

hence

$$x_{kn}^{-1}+\sum_{r=k+1}^{n-1}x_{kr}x_{rn}^{-1}+x_{kn}=0=x_{kn}+\sum_{r=k+1}^{n-1}x_{kr}x_{rn}^{-1}+x_{kn}^{-1},\quad k<n,$$

and

$$x_{kn}^{-1}=-x_{kn}-\sum_{r=k+1}^{n-1}x_{kr}x_{rn}^{-1}=-x_{kn}-\sum_{r=k+1}^{n-1}x_{kr}^{-1}x_{rn}. \tag{13}$$

We can write also

$$x_{kn}^{-1}=-\sum_{r=k+1}^{n}x_{kr}x_{rn}^{-1}=-\sum_{r=k}^{n-1}x_{kr}^{-1}x_{rn}. \tag{14}$$

There is also the explicit formula for x_{kn}^{-1} (see [8] formula (4.4)) $x_{kk+1}^{-1}=-x_{kk+1}$

$$x_{kn}^{-1}=-x_{kn}+\sum_{r=1}^{n-k-1}(-1)^{r+1}\sum_{k\leq i_1<i_2<\ldots<i_r\leq n}x_{ki_1}x_{i_1i_2}\ldots x_{i_rn},\quad k<n-1. \tag{15}$$

Remark 15 *Using (15) we see that* x_{kn}^{-1} *depends only on* x_{rs} *with* $k\leq r<s\leq n$.

Using (14) we have

$$x_{kn}+x^{-1}_{kn}=-\sum_{r=k+1}^{n-1}x_{kr}x^{-1}_{rn},\quad x_{kn}-x^{-1}_{kn}=2x_{kn}-\sum_{r=k+1}^{n-1}x_{kr}x^{-1}_{rn}. \tag{16}$$

Let us denote

$$w_{kn}:=w_{kn}(x):=(x_{kn}+x^{-1}_{kn})(x_{kn}-x^{-1}_{kn}). \tag{17}$$

Using (1) we get

$$\Delta(x)=\frac{d\mu_b(x)}{d\mu_b(x^{-1})}=\exp\left[-\sum_{k<n}b_{kn}\left(x^2_{kn}-(x^{-1}_{kn})^2\right)\right]=\exp\left[-\sum_{k<n}b_{kn}w_{kn}(x)\right]. \tag{18}$$

$$-\ln\Delta(x)=\sum_{k<n}b_{kn}\left[x^2_{kn}-(x^{-1}_{kn})^2\right]=\sum_{k<n}b_{kn}(x_{kn}+x^{-1}_{kn})(x_{kn}-x^{-1}_{kn})$$

$$\sum_{k<n}b_{kn}(x_{kn}+x^{-1}_{kn})[2x_{kn}-(x_{kn}+x^{-1}_{kn})]=\sum_{k<n}b_{kn}w_{kn}(x).$$

To study the action of the operators $A^R_{kn}=\sum_{r=1}^{k-1}x_{rk}D_{rn}+D_{kn}$ on the function $\ln\Delta(x)$ we need to know the action of D_{pq} on x^{-1}_{kn}.

Lemma 16 *We have*

$$[D_{pq},x^{-1}_{kn}]=\begin{cases}-x^{-1}_{kp}x^{-1}_{qn}, & \textit{if } k\le p<q\le n,\\ 0, & \textit{otherwise}.\end{cases} \tag{19}$$

PROOF. We prove (19) by induction in $p: k\le p<q\le n$. For $p=k$ using (16) we have

$$[D_{kq},x^{-1}_{kn}]=-[D_{kq},x_{kn}+\sum_{r=k+1}^{n-1}x_{kr}x^{-1}_{rn}]=-[D_{kq},x_{kq}x^{-1}_{qn}]=-x^{-1}_{qn}=-x^{-1}_{kk}x^{-1}_{qn},$$

so (19) holds for $p=k$.

Let us suppose that (19) holds for all (p,q) with $k\le p<s\le n,\ k\le p<q\le n$. We prove that than (19) holds also for $(s,q)\ :\ s<q\le n$. Indeed we have

$$[D_{sq},x^{-1}_{kn}]=-[D_{sq},x^{-1}_{kn}+\sum_{r=k+1}^{n-1}x_{kr}x^{-1}_{rn}]=-\sum_{r=k+1}^{s}x_{kr}[D_{sq},x^{-1}_{rn}]$$

$$=\sum_{r=k+1}^{s}x_{kr}x^{-1}_{rs}x^{-1}_{qn}\overset{(13)}{=}x^{-1}_{ks}x^{-1}_{qn}.$$

□

Using (19) we get

$$[D_{pq}, x_{kn} + x_{kn}^{-1}] = \begin{cases} -x_{kp}^{-1} x_{qn}^{-1}, & \text{if } k \leq p < q \leq n,\ (p,q) \neq (k,n) \\ 0, & \text{otherwise}. \end{cases} \tag{20}$$

Using (20) we have

$$[D_{pq}, (x_{kn}+x_{kn}^{-1})(x_{kn}-x_{kn}^{-1})] = \begin{cases} 2\, x_{kp}^{-1} x_{qn}^{-1} x_{kn}^{-1}, & \text{if } k \leq p < q \leq n,\ (p,q) \neq (k,n) \\ 2(x_{kn} + x_{kn}^{-1}), & \text{if } \quad (p,q) = (k,n) \\ 0, & \text{otherwise}. \end{cases} \tag{21}$$

Indeed, if $k \leq p < q \leq n$, $(p,q) \neq (k,n)$ we have

$$[D_{pq}, (x_{kn} + x_{kn}^{-1})(x_{kn} - x_{kn}^{-1})] = [D_{pq}, (x_{kn} + x_{kn}^{-1})(2x_{kn} - (x_{kn} + x_{kn}^{-1}))]$$

$$= [D_{pq}, (x_{kn} + x_{kn}^{-1})](2x_{kn} - (x_{kn} + x_{kn}^{-1})) - (x_{kn} + x_{kn}^{-1})[D_{pq}, (x_{kn} + x_{kn}^{-1})] =$$

$$-2x_{kn}^{-1}[D_{pq}, (x_{kn} + x_{kn}^{-1})] \overset{(20)}{=} 2x_{kp}^{-1} x_{qn}^{-1} x_{kn}^{-1}.$$

Lemma 17 *We have*

$$[A_{mm+1}^R,\, w_{kn}] = \begin{cases} 0, & \textit{if } \ k < n \leq m \\ 2x_{km} x_{km+1} & \textit{if } \ n = m+1,\ 1 \leq k \leq m-1 \\ 0, & \textit{if } \ 1 \leq k \leq m-1,\ m+1 < n \\ 2x_{mn}^{-1} x_{m+1n}^{-1}, & \textit{if } \ k = m,\ n \geq m+2 \\ 0, & \textit{if } \ \ m+1 \leq k < n. \end{cases} \tag{22}$$

hence

$$-\,[A_{mm+1}^R,\, \ln \Delta] = 2 \sum_{r=1}^{m-1} b_{rm+1} x_{rm} x_{rm+1} + 2 \sum_{n=m+2}^{\infty} b_{mn} x_{mn}^{-1} x_{m+1n}^{-1}. \tag{23}$$

PROOF. Since

$$A_{mm+1}^R = \sum_{r=1}^{m-1} x_{rm} D_{rm+1} + D_{mm+1}$$

and $w_{kn},\ k < n \leq m$ do not depend on $x_{rm+1},\ 1 \leq r \leq m+1$ we conclude that $[A_{mm+1}^R,\, w_{kn}] = 0$ for $k < n \leq m$ and $m+1 \leq k < n$.

Let $n = m+1$, since $[D_{rm+1},\, w_{km+1}] = 0$ for $1 \le r < k$ we get

$$[A^R_{mm+1},\, w_{km+1}] = \sum_{r=k}^{m-1} x_{rm}[D_{rm+1},\, w_{km+1}] + [D_{mm+1},\, w_{km+1}] =$$

$$2\left(x_{km}(x_{km+1} + x^{-1}_{km+1}) + \sum_{r=k+1}^{m-1} x_{rm}x^{-1}_{kr}x^{-1}_{km+1} + x^{-1}_{km}x^{-1}_{km+1}\right) =$$

$$2\left(x_{km}x_{km+1} + \left(x_{km} + \sum_{r=k+1}^{m-1} x^{-1}_{kr}x_{rm} + x^{-1}_{km}\right)x^{-1}_{km+1}\right) \overset{(13)}{=} 2x_{km}x_{km+1}.$$

Similarly, for $1 \le k \le m-1,\ m+1 < n$ we get

$$[A^R_{mm+1},\, w_{kn}] = \sum_{r=k}^{m-1} x_{rm}[D_{rm+1},\, w_{kn}] + [D_{mm+1},\, w_{kn}] =$$

$$2\left(x_{km}x^{-1}_{m+1n} + \sum_{r=k+1}^{m-1} x_{rm}x^{-1}_{kr}x^{-1}_{m+1n} + x^{-1}_{km}x^{-1}_{m+1n}\right)$$

$$2\left(x_{km} + \sum_{r=k+1}^{m-1} x_{rm}x^{-1}_{kr} + x^{-1}_{km}\right)x^{-1}_{m+1n} \overset{(13)}{=} 0.$$

Finally if $k = m$ and $n \ge m+2$ we have as before

$$[A^R_{mm+1},\, w_{mn}] = [D_{mm+1},\, w_{mn}] \overset{(21)}{=} 2x^{-1}_{mn}x^{-1}_{m+1n}.$$

□

We consider the action of A^R_{mm+1} on $\ln\Delta$.

Let $m = 2$. Since

$$[A^R_{23},\, w_{13}] = 2b_{13}x_{12}x_{13},\ \ [A^R_{23},\, w_{1n}] = 0,\ \ n \ge 4,\ \ [A^R_{23},\, w_{kn}] = 0,\ \ 3 \le k < n,$$

we have

$$-[A^R_{23},\, \ln\Delta] = 2b_{13}x_{12}x_{13} + 2\sum_{n=4}^{\infty} b_{2n}x^{-1}_{2n}x^{-1}_{3n},$$

hence

$$\begin{aligned} -[A^R_{12}, [A^R_{23}, \ln\Delta]] &= 2b_{13}x_{13}, \\ -[A^R_{13}, [A^R_{23}, \ln\Delta]] &= 2b_{13}x_{12}. \end{aligned}$$

The last two equations gives us $x_{12}, x_{13}\, \eta\, \mathfrak{A}$.

Let $m = 3$. Since

$$[A^R_{34},\, w_{13}] = 0,\ \ [A^R_{34},\, w_{14}] = 2x_{13}x_{14},\ \ [A^R_{34},\, w_{24}] = 2x_{23}x_{24},$$

$$[A_{34}^R,\, w_{1n}] = [A_{34}^R,\, w_{1n}] = 0,\ [A_{34}^R,\, w_{3n}] = b_{3n}x_{3n}^{-1}x_{4n}^{-1}, n \geq 5,$$
$$[A_{34}^R,\, w_{kn}] = 0,\ 4 \leq k < n,$$

we have

$$-[A_{34}^R,\, \ln\Delta] = 2b_{14}x_{13}x_{14} + 2b_{24}x_{23}x_{24} + 2\sum_{n=5}^{\infty} b_{3n}x_{3n}^{-1}x_{4n}^{-1},$$

hence

$$-[A_{23}^R, [A_{34}^R,\, \ln\Delta]] = 2b_{14}x_{12}x_{14} + 2b_{24}x_{24}$$
$$-[A_{12}^R[A_{23}^R, [A_{34}^R,\, \ln\Delta]]] = 2b_{14}x_{14},$$
$$-[A_{24}^R, [A_{34}^R,\, \ln\Delta]] = 2[x_{12}D_{14}+D_{24}, b_{14}x_{13}x_{14}+b_{24}x_{23}x_{24}] = 2b_{14}x_{12}x_{13}+2b_{24}x_{23},$$

Since $x_{12}, x_{13}\, \eta\, \mathfrak{A}$ from the latter equation we conclude that $x_{23}\, \eta\, \mathfrak{A}$. The previous equation gives us $x_{14}\, \eta\, \mathfrak{A}$ and the equation before gives $x_{24}\, \eta\, \mathfrak{A}$. Finally we conclude that $x_{14}, x_{24}, x_{23}\, \eta\, \mathfrak{A}$.

Let us suppose that we have obtained the variables $x_{rm}, 1 \leq r \leq m-2$ and $x_{m-2,m-1}$. We prove that we can obtain the following variables $x_{rm+1}, 1 \leq r \leq m-1$ and x_{m-1m}.

Indeed we calculate the action of the following sequence of operators on the result: $A_{m-1,m}^R$, $A_{m-2,m-1}^R$ etc. till A_{12}^R. We obtain

$$-[A_{m-1,m}^R, [A_{mm+1}^R,\, \ln\Delta]] = 2\left(\sum_{r=1}^{m-2} b_{r,m+1}x_{r-1,m}x_{r,m+1} + b_{m-1,m+1}x_{m-1,m+1}\right),$$

$$-[A_{m-2,m-1}^R, [A_{m-1,m}^R, [A_{mm+1}^R,\, \ln\Delta]]]$$
$$= 2\left(\sum_{r=1}^{m-3} b_{r,m+1}x_{r-2,m}x_{r,m+1} + b_{m-2,m+1}x_{m-2,m+1}\right),$$
$$-[A_{m-s,m-s+1}^R, [A_{m-s+1,m-s+2}^R, ...[A_{m-1,m}^R[A_{mm+1}^R,\, \ln\Delta]]...]]$$
$$= 2\left(\sum_{r=1}^{m-s-1} b_{r,m+1}x_{r,m-s}x_{r,m+1} + b_{m-s,m+1}x_{m-s,m+1}\right),\ 1 \leq s \leq m,$$
$$-[A_{34}^R, ...[A_{mm+1}^R,\, \ln\Delta]...] = 2(b_{1,m+1}x_{13}x_{1,m+1}+b_{2,m+1}x_{23}x_{2,m+1}+b_{3,m+1}x_{3,m+1}),$$
$$-[A_{23}^R, [A_{34}^R, ...[A_{mm+1}^R,\, \ln\Delta]...]] = 2(b_{1,m+1}x_{12}x_{1,m+1} + b_{2,m+1}x_{2,m+1}),$$
$$-[A_{12}^R, [A_{23}^R, [A_{34}^R, ...[A_{mm+1}^R,\, \ln\Delta]...]]] = 2b_{1,m+1}x_{1,m+1}.$$

From the latter equation we conclude that $x_{1,m+1}\ \eta\ \mathfrak{A}$. The last but one equation gives us $x_{2,m+1}\ \eta\ \mathfrak{A}$ (since $x_{12}, x_{1,m+1}\ \eta\ \mathfrak{A}$) etc. i.e : $x_{rm+1}\ \eta\ \mathfrak{A},\ 1 \leq r \leq m-1$.

$$-[A_{m-1m+1}^R, [A_{mm+1}^R,\, \ln\Delta]] = [\sum_{r=1}^{m-2} x_{rm-1}D_{rm+1}+D_{m-1m+1}, 2\sum_{r=1}^{m-1} b_{rm+1}x_{rm}x_{rm+1}] =$$

$$2\sum_{r=1}^{m-2} b_{rm+1}x_{rm-1}x_{rm} + b_{m-1,m+1}x_{m-1,m},$$

since $x_{rm-1}, x_{rm}\ \eta\ \mathfrak{A}$ for $1 \le r \le m-2$ hence $x_{m-1,m}\ \eta\ \mathfrak{A}$. $\Box$

To be sure that all this argument works we should prove that all involved operators are affiliated to the von Neumann algebra M'_ϕ defined by (7). For example if A^R_{23} and Δ (and hence $\ln\Delta$) are affiliated to the von Neumann algebra M'_ϕ, why the operator $[A^R_{23}, \ln\Delta]$ is also affiliated. In general, why the operators $[A^R_{12}, [A^R_{23}, [A^R_{34}, ...[A^R_{mm+1}, \ln\Delta]...]]]$ are affiliated?

Remark 18 *In general we do not know whether the commutator $[A, B]$ of two operators A and B affiliated to the von Neumann algebra is also affiliated.*

This is the reason, why we use another approach to prove that the algebra M_ϕ is trivial.

4 The von Neumann algebra M_ϕ is trivial

Since $M_\phi = (\Delta^{it},\, T^{R,b}_s \mid t \in \mathbb{R},\, s \in B^{\mathbb{N}}_0)'$ (see (7)) it is sufficient to prove that the set of operators

$$(\Delta^{is},\, T^{R,b}_t \mid s \in \mathbb{R},\, t \in B^{\mathbb{N}}_0) \subset M'_\phi$$

is irreducible.

Idea of the proof. *We show that the von Neumann subalgebra in the algebra M'_ϕ, generated by the following operators*

$$(\{T^R_{t_n}, \{T^R_{t_{n-1}}, ...\{T^R_{t_1}, \Delta^{is}\}...\}\} \mid s \in \mathbb{R},\ t_1, ..., t_n \in B^{\mathbb{N}}_0), \qquad (24)$$

where $\{a, b\} := aba^{-1}b^{-1}$ is **the maximal abelian subalgebra**. *More precisely we prove that this subalgebra contains all functions* $\exp(isx_{kn})$, $k < n,\ s \in \mathbb{R}$.

To prove the irreducibility of the algebra M'_ϕ (see proof of the Lemma 14) we observe that if an bounded operator commute with all $\exp(isx_{kn}),\ k < n,\ s \in \mathbb{R}$ then this operator itself is an operator of multiplication by some essentially bounded function $A = a(x)$. Commutation relation $[T^{R,b}_t, A] = 0$ for all $t \in B^{\mathbb{N}}_0$ gives us $a(xt) = a(x) \,\mathrm{mod}\mu_b$ for all t. Since the measure μ_b is $B^{\mathbb{N}}_0-$**right ergodic** we conclude that A is trivial i.e. $A = a(x) = CI$.

We note that expressions in (24) are the "right" analog of the left hand side of the expressions (10) and (11)

$$[A^R_{13}, [A^R_{23}, \ln\Delta]] = 2b_{13}x_{12},$$

$$[A^R_{12}, [A^R_{23}, \ln\Delta]] = 2b_{13}x_{13},$$

involving generators A^R_{kn}. In general, if we have two subgroups of unitary operators $U(t)$ and $V(s)$ with the generators A and B, to obtain the commutator $[iA, iB]$ it is sufficient to differentiate the following expression $U(t)V(s)U(-t)$:

$$\frac{\partial}{\partial t}\frac{\partial}{\partial s}U(t)V(s)U(-t)\mid_{t=s=0}=[iA, iB].$$

Indeed we have

$$\frac{\partial}{\partial s}U(t)V(s)U(-t) = U(t)iBV(s)U(-t), \quad \frac{\partial}{\partial t}U(t)iBV(s)U(-t)\mid_{t=s=0}=$$

$$(iAU(t)iBV(s)U(-t) - U(t)iBV(s)iAU(-t))\mid_{t=s=0}=[iA, iB].$$

We show that more convenient analog of the commutator $[iA, iB]$ is *commutator* (in the group sence) of two one-parameter groups

$$\{U(t), V(s)\} := U(t)V(s)U(t)^{-1}V(s)^{-1} = U(t)V(s)U(-t)V(-s).$$

Lemma 19 *For the operator g of multiplication on the function $g : f(x) \mapsto g(x)f(x)$ in the space $H_b = L_2(B^{\mathbb{N}}, d\mu_b)$ we have*

$$T^R_t g(x) T^R_{t^{-1}} = g(xt),\ t \in B^{\mathbb{N}}_0.$$

PROOF. We have

$$f(x) \overset{g(x)T^R_{t^{-1}}}{\mapsto} g(x)\left(\frac{d\mu(xt^{-1})}{d\mu(x)}\right)^{1/2} f(xt^{-1}) \overset{T^R_t}{\mapsto}$$

$$\left(\frac{d\mu(xt)}{d\mu(x)}\right)^{1/2} g(xt)\left(\frac{d\mu(x)}{d\mu(xt)}\right)^{1/2} f(x) = g(xt)f(x).$$

□

Using the lemma we have

$$T^R_t \Delta^{is}(x) T^R_{t^{-1}} = \Delta^{is}(xt).$$

Using (18) we have

$$\Delta^{is}(x) = \exp\left(-is\sum_{k+1<n} b_{kn}(x_{kn} + x^{-1}_{kn})[2x_{kn} - (x_{kn} + x^{-1}_{kn})]\right) =$$

$$\exp\left(-is\sum_{k+1<n} b_{kn}w_{kn}(x)\right), \tag{25}$$

where $w_{kn}(x) = (x_{kn} + x_{kn}^{-1})[2x_{kn} - (x_{kn} + x_{kn}^{-1})]$ (see (17)).

We would like **to obtain the functions** $\exp(isx_{kn})$ using the expressions (24). To simplify the situation we consider firstly the projections of all considered object: the measure $\mu_b^{(k)}$, the generators $A_{kn}^{R,(k)}$, operator $\Delta_{(k)}$ algebra $M^{(k)} := (M'_\phi)^{(k)}$ etc. on the following subspace $X^{(k)},\ k \geq 2$ of the space $B^{\mathbb{N}}$:

$$X^{(2)} = \left(\begin{smallmatrix} 1 & x_{12} & x_{13} & \dots & x_{1n} & \dots \\ 0 & 1 & x_{23} & \dots & x_{2n} & \dots \end{smallmatrix}\right), \quad X^{(3)} = \left(\begin{smallmatrix} 1 & x_{12} & x_{13} & x_{14} & \dots & x_{1n} & \dots \\ 0 & 1 & x_{23} & x_{14} & \dots & x_{2n} & \dots \\ 0 & 0 & 1 & x_{34} & \dots & x_{3n} & \dots \end{smallmatrix}\right), \text{ etc.}$$

Note that

$$\left(\begin{smallmatrix} 1 & x_{12} & x_{13} & \dots & x_{1n} & \dots \\ 0 & 1 & x_{23} & \dots & x_{2n} & \dots \end{smallmatrix}\right)^{-1} = \left(\begin{smallmatrix} 1 & -x_{12} & -x_{13}+x_{12}x_{23} & \dots & -x_{1n}+x_{12}x_{2n} & \dots \\ 0 & 1 & -x_{23} & \dots & -x_{2n} & \dots \end{smallmatrix}\right). \tag{26}$$

We have for the corresponding projections on $X^{(2)}$:

$$A_{1n}^R = D_{1n}, \quad A_{2n}^R = x_{12}D_{1n} + D_{2n}, \quad A_{kn}^{R,(2)} = x_{1k}D_{1n} + x_{2k}D_{2n},\ 2 < k < n,$$

$$w_{1n}(x) = (x_{1n} + x_{1n}^{-1})(x_{1n} - x_{1n}^{-1}) = x_{12}x_{2n}(2x_{1n} - x_{12}x_{2n}),\ w_{2n}(x) = 0,$$

hence

$$\Delta_{(2)}^{is}(x) := \exp\left(-is\sum_{k=3}^{\infty} b_{1n}w_{1n}(x)\right) = \exp\left(-is\sum_{k=3}^{\infty} b_{1n}x_{12}x_{2n}(2x_{1n} - x_{12}x_{2n})\right).$$

Let us denote by

$$E_{kn}(t) := I + tE_{kn}, \quad T_{kn}(t) = T^R_{E_{kn}(t)},\ k < n,\ t \in \mathbb{R} \tag{27}$$

the corresponding one-parameter subgroups. We have

$$\left(\begin{smallmatrix} x_{12} & x_{1m} \\ 1 & x_{2m} \end{smallmatrix}\right) \overset{E_{2m}(t)}{\longmapsto} \left(\begin{smallmatrix} x_{12} & x_{1m}+tx_{12} \\ 1 & x_{2m}+t \end{smallmatrix}\right),\ w_{1n}(xE_{2m}(t)) = \left\{\begin{smallmatrix} w_{1n}(x) & \text{if } n \neq m \\ w_{1m}(xE_{2m}(t)) & \text{if } n=m \end{smallmatrix}\right.$$

so using Lemma 19 we get

$$\{T_{2m}(t), \Delta_{(2)}^{is}(x)\} = T_{2m}(t)\Delta_{(2)}^{is}(x)T_{2m}(-t)\Delta_{(2)}^{-is}(x) = \Delta_{(2)}^{is}(xE_{2m}(t))\Delta_{(2)}^{-is}(x) =$$

$$\exp\left(-is\left[\sum_{k=3, k\neq m}^{\infty} b_{1n}w_{1n}(x) + b_{1m}w_{1m}(xE_{2m}(t))\right]\right)\exp\left(is\sum_{k=3}^{\infty} b_{1n}w_{1n}(x)\right) =$$

$$\exp\left(-isb_{1m}[w_{1m}(xE_{2m}(t)) - w_{1m}(x)]\right) = \exp\left(isb_{1m}(2tx_{12}x_{1m} + t^2x_{12}^2)\right),$$

since

$$w_{1m}(xE_{2m}(t)) - w_{1m}(x) = x_{12}(x_{2m}+t)[2(x_{1m}+tx_{12}) - x_{12}(x_{2m}+t)] -$$

$$x_{12}x_{2m}(2x_{1m} - x_{12}x_{2m}) = x_{12}[tx_{12}x_{2m} + t(2x_{1m} - x_{12}x_{2m}) + t^2x_{12}] = 2tx_{12}x_{1m} + t^2x_{12}^2.$$

Let us denote

$$\phi_{t,s}(x) := \{T_{2m}(t), \Delta_{(2)}^{is}(x)\} = \exp\left(isb_{1m}(2tx_{12}x_{1m} + t^2x_{12}^2)\right). \tag{28}$$

Using Lemma 19 we get

$$\{T_{1m}(t_1),\{T_{2m}(t),\Delta^{is}_{(2)}(x)\}\}=\{T_{1m}(t_1),\phi_{t,s}(x)\}=$$

$$T_{1m}(t_1)\phi_{t,s}(x)T_{1m}(-t_1)(\phi_{t,s}(x))^{-1}=\phi_{t,s}(xE_{1m}(t_1))(\phi_{t,s}(x))^{-1}=$$
$$\exp\left[isb_{1m}(2tx_{12}(x_{1m}+t_1)+t^2x_{12}^2)-isb_{1m}(2tx_{12}x_{1m}+t^2x_{12}^2)\right]=$$
$$\exp\left(isb_{1m}x_{12}2tt_1\right).$$

Finally we get for $X^{(2)}$

$$\exp(isx_{12})\in M^{(2)}:=(M'_\phi)^{(2)}.$$

Using (28) we conclude that

$$\exp(isx_{12}x_{1m})\in M^{(2)}.$$

Applying again $T_{12}(t)$ and $T_{1m}(t)$ we get

$$\{T_{12}(t),\exp(isx_{12}x_{1m})\}=T_{12}(t)\exp(isx_{12}x_{1m})T_{12}(-t)\exp(-isx_{12}x_{1m})=$$

$$\exp(is(x_{12}+t)x_{1m}-isx_{12}x_{1m})=\exp(istx_{12}),$$
$$\{T_{1m}(t),\exp(isx_{12}x_{1m})\}=T_{1m}(t)\exp(isx_{12}x_{1m})T_{1m}(-t),\exp(-isx_{12}x_{1m})=$$
$$\exp(isx_{12}(x_{1m}+t)-isx_{12}x_{1m})=\exp(istx_{1m}).$$

At last we conclude that for $X^{(2)}$ we have $\exp(isx_{12}),\ \exp(isx_{1m})\in M^{(2)}$ in particular

$$\exp(isx_{12}),\ \exp(isx_{13})\in M^{(2)}. \tag{29}$$

For $X^{(3)}$ and the corresponding projections we have

$$\begin{pmatrix}1&x_{12}&x_{13}&x_{14}&\dots&x_{1n}&\dots\\0&1&x_{23}&x_{14}&\dots&x_{2n}&\dots\\0&0&1&x_{34}&\dots&x_{3n}&\dots\end{pmatrix}^{-1}=$$

$$\begin{pmatrix}1&-x_{12}&-x_{13}+x_{12}x_{23}&-x_{14}+x_{12}x_{24}+x_{13}x_{34}+x_{12}x_{23}x_{34}&\dots&-x_{1n}+x_{12}x_{2n}+x_{13}x_{3n}+x_{12}x_{23}x_{3n}&\dots\\0&1&-x_{23}&-x_{24}+x_{23}x_{34}&\dots&-x_{2n}+x_{23}x_{3n}&\dots\\0&0&1&-x_{34}&\dots&x_{3n}&\dots\end{pmatrix}=$$
$$\begin{pmatrix}1&-x_{12}&-x_{13}-x_{12}^{-1}x_{23}&-x_{14}-x_{12}^{-1}x_{24}-x_{13}^{-1}x_{34}&\dots&-x_{1n}-x_{12}^{-1}x_{2n}-x_{13}^{-1}x_{3n}&\dots\\0&1&-x_{23}&-x_{24}-x_{23}^{-1}x_{34}&\dots&-x_{2n}-x_{23}^{-1}x_{3n}&\dots\\0&0&1&-x_{34}&\dots&-x_{3n}&\dots\end{pmatrix}, \tag{30}$$

$$A^R_{1n}=D_{1n},\quad A^R_{2n}=x_{12}D_{1n}+D_{2n},\quad A^R_{3n}=x_{13}D_{1n}+x_{23}D_{2n}+D_{3n},\ 3<n.$$

We have

$$\Delta^{is}_{(3)}(x)=\exp\left(-is\left[\sum_{n=3}^{\infty}b_{1n}w_{1n}(x)+\sum_{n=4}^{\infty}b_{2n}w_{2n}(x)\right]\right)=$$

$$\exp\left(-is\left[\sum_{n=3}^{\infty}b_{1n}(x_{1n}+x_{1n}^{-1})[2x_{1n}-(x_{1n}+x_{1n}^{-1})]\right]\right)\times$$
$$\exp\left(-is\left[\sum_{n=4}^{\infty}b_{2n}(x_{2n}+x_{2n}^{-1})[2x_{2n}-(x_{2n}+x_{2n}^{-1})\right]\right).$$

By the same procedure as in the case of the space $X^{(2)}$ we can obtain that

$$\exp(isx_{12}),\ \exp(isx_{13}) \in M^{(3)}. \tag{31}$$

We show that

$$\{T_{34}(t), \Delta^{is}_{(3)}(x)\} = \exp\left(is\left[b_{14}(2tx_{13}x_{14}+t^2x_{13}^2)+b_{24}(2tx_{23}x_{24}+t^2x_{23}^2)\right]\right). \tag{32}$$

(compare with (28)). Indeed we have

$$\{T_{34}(t), \Delta^{is}_{(3)}(x)\} = T_{34}(t)\Delta^{is}_{(3)}(x)T_{34}(-t)\Delta^{-is}_{(3)}(x) =$$

$$\Delta^{is}_{(3)}(xE_{34}(t))\Delta^{-is}_{(3)}(x) =$$

$$\exp\left(-is\left(b_{14}[w_{14}(xE_{34}(t))-w_{14}(x)]+b_{24}[w_{24}(xE_{34}(t))-w_{24}(x)]\right)\right),$$

which implies (32), since

$$w_{14}(x) = (x_{14}+x_{14}^{-1})[2x_{14}-(x_{14}+x_{14}^{-1})] = -(x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34})[2x_{14}+x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34}],$$

and

$$w_{14}(xE_{34}(t)) - w_{14}(x) =$$

$$-[x_{12}^{-1}(x_{24}+tx_{23})+x_{13}^{-1}(x_{34}+t)][2(x_{14}+tx_{13})+x_{12}^{-1}(x_{24}+tx_{23})+x_{13}^{-1}(x_{34}+t)]$$

$$+(x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34})[2x_{14}+x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34}] =$$

$$-t\left[(x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34})(2x_{13}+x_{12}^{-1}x_{23}+x_{13}^{-1})+(x_{12}^{-1}x_{23}+x_{13}^{-1})(2x_{14}+x_{12}^{-1}x_{24}+x_{13}^{-1}x_{34})\right]$$

$$-t^2(x_{12}^{-1}x_{23}+x_{13}^{-1})(2x_{13}+x_{12}^{-1}x_{23}+x_{13}^{-1}) = -t[-(x_{14}+x_{14}^{-1})x_{13}-x_{13}(x_{14}+x_{14}^{-1})]+t^2x_{13}x_{13} =$$

$$2tx_{13}x_{14}+t^2x_{13}^2.$$

Using (31) and (32) we get

$$\phi^{(3)}_{t,s}(x) := \exp\left(is\left[b_{14}2tx_{13}x_{14}+b_{24}(2tx_{23}x_{24}+t^2x_{23}^2)\right]\right) \in M^{(3)},$$

hence

$$\{T_{13}(t_1), \phi^{(3)}_{t,s}(x)\} = T_{13}(t_1)\phi^{(3)}_{t,s}(x)T_{13}(-t_1)(\phi^{(3)}_{t,s}(x))^{-1} = \exp\left(istt_1b_{14}2tx_{14}\right),$$

so $\exp(isx_{14}) \in M^{(3)}$ and $\exp[isb_{24}(2tx_{23}x_{24}+t^2x_{23}^2)] \in M^{(3)}$. Similarly we get

$$\{T_{24}(t_1), \exp[isb_{24}(2tx_{23}x_{24}+t^2x_{23}^2)]\} = \exp(isb_{24}tt_1x_{23}),$$

so $\exp(isx_{23}), \exp(isx_{23}x_{24}) \in M^{(3)}$. At last we get

$$\{T_{24}(t_1), \exp(isx_{23}x_{24})\} = \exp(ist_1x_{24}).$$

Finally we can obtain $\exp(isx_{kn})$ in the following order on the **first step**:

$$\exp(isx_{12}),\ \exp(isx_{13});$$

on the **second step**:

$$\exp(isx_{14}),\ \exp(isx_{23}),\ \exp(isx_{24})\in M^{(3)},$$

or symbolically in the following **order**:

$$\begin{pmatrix} 1 & x_{12} & x_{13} & x_{14}\\ 0 & 1 & x_{23} & x_{24}\\ 0 & 0 & 1 & \end{pmatrix},\quad \begin{pmatrix} 0 & 1_1 & 2_1 & 1_2\\ 0 & 0 & 2_2 & 3_2\\ 0 & 0 & 0 & \end{pmatrix}.$$

In general we get **the order**

$$\begin{pmatrix} 1 & x_{12} & x_{13} & x_{14} & x_{15} & x_{16} & x_{17}\\ 0 & 1 & x_{23} & x_{24} & x_{25} & x_{26} & x_{27}\\ 0 & 0 & 1 & x_{34} & x_{35} & x_{36} & x_{37}\\ 0 & 0 & 0 & 1 & x_{45} & x_{46} & x_{47}\\ 0 & 0 & 0 & 0 & 1 & x_{56} & x_{57}\\ 0 & 0 & 0 & 0 & 0 & 1 & \end{pmatrix},\quad \begin{pmatrix} 0 & 1_1 & 2_1 & 1_2 & 1_3 & 1_4 & 1_5\\ 0 & 0 & 2_2 & 3_2 & 2_3 & 2_4 & 2_5\\ 0 & 0 & 0 & 3_3 & 4_3 & 3_4 & 3_5\\ 0 & 0 & 0 & 0 & 4_4 & 5_4 & 4_5\\ 0 & 0 & 0 & 0 & 0 & 5_5 & \end{pmatrix}. \tag{33}$$

This **order is right in the general case** (without any projections on $X^{(k)}$). To obtain $\exp(isx_{12})$ and $\exp(isx_{13})$ on the **first step** we get by Lemma 19

$$\{T_{23}(t),\Delta^{is}(x)\}=T_{23}(t)\Delta^{is}(x)T_{23}(-t)\Delta^{-is}(x)=\Delta^{is}(xE_{23}(t))\Delta^{-is}(x)=$$

$$\exp\left\{-is\left(\sum_{n=3}^{\infty}b_{1n}[w_{1n}(xE_{23}(t))-w_{1n}(x)]+\sum_{n=4}^{\infty}b_{2n}[w_{2n}(xE_{23}(t))-w_{2n}(x)]\right)\right\}. \tag{34}$$

Now we shall calculate $w_{1n}(xE_{23}(t))-w_{1n}(x)$ and $w_{2n}(xE_{23}(t))-w_{2n}(x)$. We have by (16)

$$x_{1n}+x^{-1}_{1n}=-\sum_{r=2}^{n-1}x_{1r}x^{-1}_{rn},\quad x_{2n}+x^{-1}_{2n}=-\sum_{r=3}^{n-1}x_{2r}x^{-1}_{rn}$$

so we conclude that for $n>3$ holds

$$(x_{1n}+x^{-1}_{1n})^{E_{23}(t)}=-\left(\sum_{r=2}^{n-1}x_{1r}x^{-1}_{rn}\right)^{E_{23}(t)}=-\left(x_{12}x^{-1}_{2n}+x_{13}x^{-1}_{3n}+\sum_{r=4}^{n-1}x_{1r}x^{-1}_{rn}\right)^{E_{23}(t)}=$$

$$-\left(x_{12}(-x_{2n}-[x_{23}+t]x^{-1}_{3n}-\sum_{r=4}^{n-1}x_{2r}x^{-1}_{rn})+[x_{13}+tx_{12}]x^{-1}_{3n}+\sum_{r=4}^{n-1}x_{1r}x^{-1}_{rn}\right)=$$

$$-\left(\sum_{r=2}^{n-1}x_{1r}x^{-1}_{rn}-tx_{12}x^{-1}_{3n}+tx_{12}x^{-1}_{3n}\right)=x_{1n}+x^{-1}_{1n}.$$

For $n=3$ we get $x_{13}+x^{-1}_{13}=-x_{12}x^{-1}_{23}=x_{12}x_{23}$ hence

$$(x_{13}+x^{-1}_{13})^{E_{23}(t)}=(x_{12}x_{23})^{E_{23}(t)}=$$

$$x_{12}[x_{23}+t]=x_{12}x_{23}+tx_{12}=x_{13}+x^{-1}_{13}-tx^{-1}_{12}.$$

Finally we conclude that

$$(x_{1n} + x_{1n}^{-1})^{E_{23}(t)} = \begin{cases} x_{1n} + x_{1n}^{-1}, & \text{if } 3 < n, \\ x_{13} + x_{13}^{-1} + tx_{12}, & \text{if } \ n = 3 \end{cases} \tag{35}$$

and

$$(x_{1n} \pm x_{1n}^{-1})^{E_{23}(t)} = \begin{cases} x_{1n} \pm x_{1n}^{-1}, & \text{if } 3 < n, \\ x_{13} \pm x_{13}^{-1} + tx_{12}, & \text{if } \ n = 3 \end{cases} \tag{36}$$

since

$$(x_{13} - x_{13}^{-1})^{E_{23}(t)} = (2x_{13} - (x_{13} + x_{13}^{-1}))^{E_{23}(t)} = 2[x_{13} + tx_{12}] - (x_{13} + x_{13}^{-1} + tx_{12})$$

$$= x_{13} - x_{13}^{-1} + tx_{12}.$$

We have $w_{1n}(xE_{23}(t)) - w_{1n}(x) = 0$ for $n > 3$. For $n = 3$ holds

$$w_{13}(xE_{23}(t)) - w_{13}(x) = (x_{13} + x_{13}^{-1} + tx_{12})(x_{13} - x_{13}^{-1} + tx_{12}) - (x_{13} + x_{13}^{-1})(x_{13} - x_{13}^{-1})$$

$$= tx_{12}(x_{13} + x_{13}^{-1} + x_{13} - x_{13}^{-1}) + t^2x_{12}^2 = 2tx_{12}x_{13} + t^2x_{12}^2.$$

Finally

$$w_{1n}(xE_{23}(t)) - w_{1n}(x) = \begin{cases} 0, & \text{if } \quad 3 < n \\ 2tx_{12}x_{13} + t^2x_{12}^2, & \text{if } \quad n = 3. \end{cases} \tag{37}$$

For $(x_{2n} + x_{2n}^{-1})^{E_{23}(t)}$ we have

$$(x_{2n} + x_{2n}^{-1})^{E_{23}(t)} = -\left(\sum_{r=3}^{n-1} x_{2r}x_{rn}^{-1}\right)^{E_{23}(t)} = -\left(x_{23}x_{3n}^{-1} + \sum_{r=4}^{n-1} x_{2r}x_{rn}^{-1}\right)^{E_{23}(t)} =$$

$$-\left([x_{23} + t]x_{3n}^{-1} + \sum_{r=4}^{n-1} x_{2r}x_{rn}^{-1}\right) = x_{2n} + x_{2n}^{-1} - tx_{3n}^{-1}.$$

Since

$$(x_{2n} - x_{2n}^{-1})^{E_{23}(t)} = [2x_{2n} - (x_{2n} + x_{2n}^{-1})]^{E_{23}(t)} = [2x_{2n} - (x_{2n} + x_{2n}^{-1} - tx_{3n}^{-1})]$$

$$= x_{2n} - x_{2n}^{-1} + tx_{3n}^{-1}$$

we conclude that

$$(x_{2n} \pm x_{2n}^{-1})^{E_{23}(t)} = x_{2n} \pm x_{2n}^{-1} \mp tx_{3n}^{-1}. \tag{38}$$

Finally we have

$$w_{2n}(xE_{23}(t)) - w_{2n}(x) = (x_{2n} + x_{2n}^{-1} - tx_{3n}^{-1})(x_{2n} - x_{2n}^{-1} + tx_{3n}^{-1}) - (x_{2n} + x_{2n}^{-1})(x_{2n} - x_{2n}^{-1}) =$$

$$tx_{3n}^{-1}(x_{2n} + x_{2n}^{-1} + x_{2n} - x_{2n}^{-1}) - t^2(x_{3n}^{-1})^2 = 2tx_{2n}^{-1}x_{3n}^{-1} - t^2(x_{3n}^{-1})^2,$$

$$w_{2n}(xE_{23}(t)) - w_{2n}(x) = 2tx_{2n}^{-1}x_{3n}^{-1} - t^2(x_{3n}^{-1})^2. \tag{39}$$

Using (37) and (39) we get

$$w_{kn}(xE_{23}(t)) - w_{kn}(x) = \begin{cases} 2tx_{12}x_{13} + t^2x_{12}^2, & \text{if} \quad n=3,\ k=1 \\ 2tx_{2n}^{-1}x_{3n}^{-1} - t^2(x_{3n}^{-1})^2, & \text{if} \quad k=2,\ n\geq 4 \\ 0, & \text{otherwise.} \end{cases} \tag{40}$$

At last using (34) and (40) we have

$$\{T_{23}(t), \Delta^{is}(x)\} = \exp\left(-is\left[b_{13}(2tx_{12}x_{13} + t^2x_{12}^2) + \sum_{n=4}^{\infty} b_{2n}(2tx_{2n}^{-1}x_{3n}^{-1} - t^2(x_{3n}^{-1})^2)\right]\right).$$

Further we get

$$\{T_{13}(t_2)\{T_{23}(t_1), \Delta^{is}(x)\}\} = \exp\left(-isb_{13}2t_1t_2x_{12}\right). \tag{41}$$

Indeed

$$\{T_{13}(t_2)\{T_{23}(t_1), \Delta^{is}(x)\}\} =$$
$$\exp\left(-isb_{13}\left[(2t_1x_{12}[x_{13}+t_2] - t_1^2x_{12}^2) - (2t_1x_{12}x_{13} - t_1^2x_{12}^2)\right]\right)$$
$$= \exp\left(-isb_{13}2t_1t_2x_{12}\right),$$

compare with (10): $-[A_{13}^R, [A_{23}^R, \ln\Delta]] = 2b_{13}x_{12}$! We have $\exp(itx_{12}) \in M_\phi'$ and hence $\exp(itx_{12}^2) \in M_\phi'$. Using expression for $\{T_{23}(t_1), \Delta^{is}(x)\}$ we conclude that

$$M_\phi' \ni \{T_{23}(t_1), \Delta^{is}(x)\}\exp(isb_{13}t^2x_{12}^2) =$$
$$\exp\left(-is\left[b_{13}(2tx_{12}x_{13}) + \sum_{n=4}^{\infty} b_{2n}(2tx_{2n}^{-1}x_{3n}^{-1} - t^2(x_{3n}^{-1})^2)\right]\right),$$

so

$$M_\phi' \ni \{T_{12}(t_2), \{T_{23}(t_1), \Delta^{is}(x)\}\exp(isb_{13}t^2x_{12}^2)\} = \exp\left(-isb_{13}2t_1t_2x_{13}\right).$$

Compare with the expression $-[A_{12}^R, [A_{23}^R, \ln\Delta]] = 2b_{13}x_{13}$. Finally we conclude that

$$\exp(itx_{12}), \quad \exp(itx_{13}) \in M_\phi' \tag{42}$$

In general (without any projections) the following lemma holds

Lemma 20 *We have*

$$w_{kn}(xE_{mm+1}(t)) - w_{kn}(x) = \begin{cases} 2tx_{rm}x_{rm+1} + t^2x_{rm+1}^2, & \textit{if } n=m+1,\ 1\leq k\leq m-1 \\ 2tx_{mn}^{-1}x_{m+1n}^{-1} - t^2(x_{m+1n}^{-1})^2, & \textit{if } k=m,\ n\geq m+2 \\ 0, & \textit{otherwise,} \end{cases} \tag{43}$$

hence

$$\{T_{mm+1}(t), \Delta^{is}(x)\} =$$

$$\exp\left(-is\left[\sum_{r=1}^{m-1} b_{rm+1}(2tx_{rm}x_{rm+1}+t^2x^2_{rm+1}) + \sum_{n=m+2}^{\infty} b_{mn}(2tx^{-1}_{mn}x^{-1}_{m+1n} - t^2(x^{-1}_{m+1n})^2)\right]\right). \tag{44}$$

PROOF. The proof is similar to the proof of the Lemma 17. □

To obtain another functions $\exp(itx_{kn})$ in the general case we should make all the steps as it was indicated before. For example to obtain $\exp(isx_{14})$, $\exp(isx_{23})$, $\exp(isx_{24})$ we should do **the second step** i.e. consider the operators

$$\{T_{34}(t), \Delta^{is}(x)\}$$

and all necessary combinations.

To obtain $\exp(isx_{15})$, $\exp(isx_{25})$, $\exp(isx_{34})$, $\exp(isx_{34})$ we should consider the following operators

$$\{T_{45}(t), \Delta^{is}(x)\},$$

and so on. Finally we shall obtain all functions $\exp(isx_{kn})$, $k < n$.

5 Example of the measure

We show that the set $b = (b_{kn})_{k<n}$ for which

$$S^L_{kn}(b) < \infty, \quad E(b) < \infty, \quad \text{and } S^{R,L}_{kn}(b) = \infty, \ \ 1 \leq k < n,$$

where

$$S^L_{kn}(b) = \sum_{m=n+1}^{\infty} \frac{b_{km}}{b_{nm}}, \ E(b) = \sum_{k<n} \frac{S^L_{kn}(b)}{b_{kn}}, \ S^{R,L}_{kn}(b) = \sum_{m=n+1}^{\infty} \frac{b_{km}}{S^L_{nm}(b)}.$$

is not empty. Indeed let us take $b_{kn} = (a_k)^n$. We have

$$S^L_{kn}(b) = \sum_{m=n+1}^{\infty} \left(\frac{a_k}{a_n}\right)^m = \left(\frac{a_k}{a_n}\right)^{n+1} \sum_{m=0}^{\infty} \left(\frac{a_k}{a_n}\right)^m = \left(\frac{a_k}{a_n}\right)^{n+1} \frac{1}{1-\frac{a_k}{a_n}} < \infty$$

iff $a_k < a_{k+1}, \ k \in \mathbb{N}$, for example $a_k = s^k$ with $s > 1$. Further we get

$$E(b) = \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} \frac{S^L_{kn}(b)}{b_{kn}} = \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} \left(\frac{a_k}{a_n}\right)^{n+1} \frac{1}{1-\frac{a_k}{a_n}} \frac{1}{a^n_k} =$$

$$\sum_{k=1}^{\infty} a_k \sum_{n=k+1}^{\infty} \left(\frac{1}{a_n}\right)^{n+1} \frac{1}{1-\frac{a_k}{a_n}} < \sum_{k=1}^{\infty} \frac{a_k}{1-\frac{a_k}{a_{k+1}}} \sum_{n=k+1}^{\infty} \left(\frac{1}{a_n}\right)^{n+1}$$

$$< \sum_{k=1}^{\infty} \frac{a_k}{1-\frac{a_k}{a_{k+1}}} \sum_{n=k+1}^{\infty} \left(\frac{1}{a_{k+1}}\right)^{n+1} = \sum_{k=1}^{\infty} \frac{a_k}{1-\frac{a_k}{a_{k+1}}} \left(\frac{1}{a_{k+1}}\right)^{k+2} \frac{1}{1-\frac{1}{a_{k+1}}} =$$

$$\sum_{k=1}^{\infty} \frac{\frac{a_k}{a_{k+1}}}{1-\frac{a_k}{a_{k+1}}} \left(\frac{1}{a_{k+1}}\right)^{k} \frac{1}{a_{k+1}-1} < \sum_{k=1}^{\infty} \frac{\frac{a_k}{a_{k+1}}}{1-\frac{a_k}{a_{k+1}}} \left(\frac{1}{a_2}\right)^{k} \frac{1}{a_2-1}.$$

If for example $a_k = s^k$ with $s > 1$ we have

$$E(b) < \frac{\frac{1}{s}}{1-\frac{1}{s}} \sum_{k=1}^{\infty} \frac{1}{s^{k(k+1)}} \frac{1}{s^{k+1}-1} < \infty.$$

At last

$$S_{kn}^{R,L}(b) = \sum_{m=n+1}^{\infty} \frac{b_{km}}{S_{nm}^{L}(b)} = \sum_{m=n+1}^{\infty} \frac{a_k^m \left(1-\frac{a_n}{a_m}\right)}{\left(\frac{a_n}{a_m}\right)^{m+1}}$$

$$= \sum_{m=n+1}^{\infty} \left(\frac{a_k a_m}{a_n}\right)^m \left(\frac{a_m}{a_n}\right)\left(1-\frac{a_n}{a_m}\right) = \sum_{m=n+1}^{\infty} \left(\frac{a_k a_m}{a_n}\right)^m \left(\frac{a_m}{a_n}-1\right) = \infty,$$

if $\lim_m a_m = \infty$. For $a_k = s^k$ with $s > 1$ we have

$$S_{kn}^{R,L}(b) = \sum_{m=n+1}^{\infty} s^{(m+k-n)m}(s^{m-n}-1) = \infty.$$

6 Modular operator

We recall how to find the modular operator and the operator of canonical conjugation for the von Neumann algebra $\mathfrak{A}_G^{\rho}$, generated by the right regular representation ρ of a locally compact Lie group G. Let h be a right invariant Haar measure on G and

$$\rho, \lambda : G \mapsto U(L^2(G,h))$$

be the right and the left regular representations of the group G defined by

$$(\rho_t f)(x) = f(xt),\ (\lambda_t f)(x) = (dh(t^{-1}x)/dh(x))^{-1/2} f(t^{-1}x).$$

To define the right Hilbert algebra on G we can proceed as follows. Let $M(G)$ be algebra of all probability measures on G with convolution

$$(\mu * \nu)(s) =$$

We define the homomorphism

$$M(G) \ni \mu \mapsto \rho^{\mu} = \int_G \rho_t \, \mathrm{d}\mu(t) \in B(L^2(G,h)).$$

We have $\rho^\mu\rho^\nu = \rho^{\mu*\nu}$, indeed

$$\rho^\mu\rho^\nu = \int_G \rho_t \,\mathrm{d}\mu(t) \int_G \rho_s \,\mathrm{d}\nu(s) = \int_G\int_G \rho_{ts} \,\mathrm{d}\mu(t)\nu(s) = \int_G \rho_t \,\mathrm{d}(\mu * \nu)(t) = \rho^{\mu*\nu}.$$

Let us consider a subalgebra $M_h(G) := (\nu \in M(G) \mid \nu \sim h)$ of the algebra $M_h(G)$ In the case when $\mu \in M_h(G)$ we can associate with the measure μ its Rodon-Nikodim derivative $d\nu(t)/dh(t) = f(t)$. When $f \in C_0^\infty(G)$ or $f \in L^1(G)$ we can write

$$\rho^f = \int_G f(t)\rho_t dh(t),$$

hence we can replace the algebra $M_h(G)$ by its subalgebra identified with algebra of functions $C_0^\infty(G)$ or $L^1(G,h)$ with convolutions. If we replace the Haar measure h with some measure $\mu \in M_h(G)$ we obtain the isomorphic image $T^{R,\mu}$ of the right regular representation ρ in the space $L^2(G,\mu)$: $T_t^{R,\mu} = U\rho_t U^{-1}$ where $U : L^2(G,h) \mapsto L^2(G,\mu)$ defined by $(Uf)(x) = \left(\frac{dh(x)}{d\mu(x)}\right)^{1/2} f(x)$. we have

$$(T_t^{R,\mu}f)(x) = \left(\frac{d\mu(xt)}{d\mu(x)}\right)^{1/2} f(xt),$$

and

$$T^f = \int_G f(t)T_t^{R,\mu} d\mu(t).$$

We have (see [4], p.462) (we shall write T_t instead of $T_t^{R,\mu}$)

$$S(T^f) := (T^f)^* = \int_G \overline{f(t)} T_{t^{-1}} d\mu(t) = \int_G \overline{f(t)} T_{t^{-1}} \frac{d\mu(t)}{d\mu(t^{-1})} d\mu(t^{-1})$$

$$\int_G \frac{d\mu(t^{-1})}{d\mu(t)} \overline{f(t^{-1})} T_t d\mu(t).$$

Hence

$$(Sf)(t) = \frac{d\mu(t^{-1})}{d\mu(t)} \overline{f(t^{-1})}.$$

To calculate S^* we use the fact that S is antilinear so $(Sf,g) = (S^*g,f)$. We have

$$(Sf,g) = \int_G \frac{d\mu(t^{-1})}{d\mu(t)} \overline{f(t^{-1})g(t)} d\mu(t) = \int_G \overline{f(t^{-1})g(t)} d\mu(t^{-1}) =$$

$$\int_G \overline{g(t^{-1})f(t)} d\mu(t) = (S^*g,f),$$

hence $(S^*g)(t) = \overline{g(t^{-1})}$. Finally the modular operator Δ defined by $\Delta = S^*S$ has the following form $(\Delta f)(t) = \frac{d\mu(t)}{d\mu(t^{-1})} f(t)$. Indeed we have

$$f(t) \overset{S}{\mapsto} \frac{d\mu(t^{-1})}{d\mu(t)} \overline{f(t^{-1})} \overset{S^*}{\mapsto} \frac{d\mu(t)}{d\mu(t^{-1})} f(t).$$

Finally , since $J = S\Delta^{-1/2}$ (see [4] p.462) we get

$$f(t) \overset{\Delta^{-1/2}}{\mapsto} \left(\frac{d\mu(t^{-1})}{d\mu(t)}\right)^{1/2} f(t) \overset{J}{\mapsto} \frac{d\mu(t^{-1})}{d\mu(t)} \left(\frac{d\mu(t)}{d\mu(t^{-1})}\right)^{1/2} \overline{f(t^{-1})}$$

$$= \left(\frac{d\mu(t^{-1})}{d\mu(t)}\right)^{1/2} \overline{f(t^{-1})}.$$

Hence

$$(Jf)(t) = \left(\frac{d\mu(t^{-1})}{d\mu(t)}\right)^{1/2} \overline{f(t^{-1})}, \text{ and } (\Delta f)(t) = \frac{d\mu(t)}{d\mu(t^{-1})} f(t).$$

To prove that $JT_t^{R,\mu}J = T_t^{L,\mu}$ we get

$$f(t) \overset{J}{\mapsto} \left(\frac{d\mu(x^{-1})}{d\mu(x)}\right)^{1/2} \overline{f(x^{-1})} \overset{T_t^{R,\mu}}{\mapsto} \left(\frac{d\mu(xt)}{d\mu(x)}\right)^{1/2} \left(\frac{d\mu((xt)^{-1})}{d\mu(xt)}\right)^{1/2} \overline{f((xt)^{-1})} =$$

$$\left(\frac{d\mu(t^{-1}x^{-1})}{d\mu(x)}\right)^{1/2} \overline{f(t^{-1}x^{-1})} \overset{J}{\mapsto} \left(\frac{d\mu(x^{-1})}{d\mu(x)}\right)^{1/2} \left(\frac{d\mu(t^{-1}x)}{d\mu(x^{-1})}\right)^{1/2} f(t^{-1}x) =$$

$$\left(\frac{d\mu(t^{-1}x)}{d\mu(x)}\right)^{1/2} f(t^{-1}x) = (T_t^{L,\mu} f)(x).$$

Remark 21 *The representation T^{R,μ_b} is the inductive limit of the representations T^{R,μ_b^m} of the group $B(m,\mathbb{R})$ where the measure μ_b^m is the projection of the measure μ_b onto subgroup $B(m,\mathbb{R})$. Obviously μ_b^m is equivalent with the Haar measure h_m on $B(m,\mathbb{R})$.*

7 The uniqueness of the constructed factor

Let G be a solvable separable locally compact group or a connected locally compact group. Then any representation π of G in a Hilbert space generates an approximately finite-dimensional von Neumann algebra (see [3]).

Theorem 15 from V.9 p. 504 [4] (Haagerup) There exists up to isomorphism only one amenable factor of type III_1, the factor R_∞ of Araki and Woods (see [7]).

Acknowledgements. The author would like to thank the Max-Planck-Institute of Mathematics where the main part of the work was done for the hospitality. He is grateful to Prof. A.Connes for useful remarks concerning the uniqueness of the constructed factor. The partial financial support by the DFG project 436 UKR 113/87 are gratefully acknowledged.

References

[1] S. Albeverio and A. Kosyak, Group action, quasi-invariant measures and quasiregular representations of the infinite-dimensional nilpotent group, Contemporary Mathematics of AMS. 385 (2005) 259-280.

[2] S. Albeverio and A. Kosyak, Quasiregular representations of the infinite-dimensional nilpotent group, J. Funct. Anal. 236 (2006) 634-681.

[3] A. Connes, Classification of injective factors. Cases II_1, II_∞, III_λ, $\lambda \neq 1$. Ann. of Math. (2) 104 (1976), no. 1, 73–115.

[4] A. Connes, Noncommutative geometry. Academic Press, Inc., San Diego, CA, 1994.

[5] J. Dixmier, Les algèbres d'opérateurs dans l'espace hilbertien, 2nd Edition, Gauthier-Villars, Paris, 1969.

[6] I. Dynov, and A. Kosyak, Type III_1 factors generated by regular representations of infinite dimensional nilpotent group $B_0^{\mathbb{Z}}$, (in preparations).

[7] U. Haagerup, Connes' bicentralizer problem and uniqueness of the injective factor of type III_1. Acta Math. 158 No. 1-2 (1987) 95–148.

[8] A.V. Kosyak, Extension of unitary representations of inductive limits of finite-dimensional Lie groups, Rep. Math. Phys. 26 No. 2 (1988) 129–148.

[9] A.V. Kosyak, Irreducibility criterion for regular Gaussian representations of group of finite upper triangular matrices, Funktsional. Anal. i Priložhen 24 No 3, (1990) 82–83 (in Russian). (transl. in Funct. Anal. Appl. 24 No 3, (1990) 243–245 (1991)).

[10] A.V. Kosyak, Criteria for irreducibility and equivalence of regular Gaussian representations of group of finite upper triangular matrices of infinite order, Selecta Math. Soviet. 11 (1992) 241–291.

[11] A.V. Kosyak, Irreducible regular Gaussian representations of the group of the interval and the circle diffeomorphisms, J. Funct. Anal. 125 (1994) 493–547.

[12] A.V. Kosyak, Inversion-quasi-invariant Gaussian measures on the group of infinite-order upper-triangular matrices, Funct. Anal. i Priloz. 34, issue 1 (2000) 86–90.

[13] A. Kosyak and R. Zekri, Regular representations of infinite-dimensional groups and factors, I, Methods of Funct. Anal. Topology. 6, No 2 (2000) 50–59.

[14] A. Kosyak and R. Zekri, Regular representations of infinite-dimensional group $B_0^{\mathbb{Z}}$ and factors. Methods of Funct. Anal. Topology. 7, No 4 (2001) 43 – 48.

[15] A.V. Kosyak, The generalized Ismagilov conjecture for the group $B_0^{\mathbb{N}}$. II, Methods Funct. Anal. Topology. 8, No 3 (2002) 27–45.

[16] M. Takesaki, Theory of operator algebras. III. Encyclopaedia of Mathematical Sciences, 127. Operator Algebras and Non-commutative Geometry, 8. Springer-Verlag, Berlin, 2003.